

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.






Defects in the images may include (but are not limited to):






- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS




IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

the sum of the energies of all of said absorbed photons is greater than or equal to the transition energy from a ground state of said compound to said multi-photon excited state and wherein the energy of each absorbed photon is less than the transition energy between said ground state and the lowest single-photon excited state of said compound and is less than the transition energy between said multi-photon excited state and said ground state [A method according to claim 2], wherein said compound is selected from the group consisting of








and mixtures thereof, where $R = (CH_2)_{11}CH_3$.

4. (Amended) A method for preparing a compound in an electronically excited state, comprising the steps of:

a) exposing a compound having the formula D_1 -II- D_2 to radiation, wherein D_1 and D_2 are electron donor groups; and II comprises a bridge of π -conjugated bonds connecting D_1 and D_2 ; and

b) converting said compound to a multi-photon electronically excited state upon simultaneous absorption of at least two photons of said radiation by said compound, wherein the sum of the energies of all of said absorbed photons is greater than or equal to the

transition energy from a ground state of said compound to said multi-photon excited state and wherein the energy of each absorbed photon is less than the transition energy between said ground state and the lowest single-photon excited state of said compound and is less than the transition energy between said multi-photon excited state and said ground state [A method according to claim 2], wherein said compound is further defined by a formula



where D_a is selected from the group consisting of N, O, S and P;

where D_b is selected from the group consisting of N, O, S and P;

m, n, o are integers such that $0 \leq m \leq 10$, $0 \leq n \leq 10$, $0 \leq o \leq 10$; and

where:

X, Y, Z are independently selected from the group consisting of: $CR_k=CR_l$; O; S; and N- R_m ;

R_a, R_b, R_c, R_d are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{a1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl; where $0 < \alpha < 10$ and $1 < \beta < 25$, a group of aromatic rings having up to 20 carbons in the aromatic ring framework; fused aromatic rings, vinyl; allyl; 4-styryl; acroyl; methacroyl; acrylonitrile, isocyanate; isothiocyanate; epoxides; strained ring olefins; $(-CH_2)_\delta SiCl_3$; $(-CH_2)_\delta Si(OCH_2CH_3)_3$; and $(-CH_2)_\delta Si(OCH_3)_3$; where $\delta < 25$;

R_{a1}, R_{a2} , and R_{a3} are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons, a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof, or methacryloyl chloride;

$R_e, R_f, R_g, R_h, R_i, R_j, R_k, R_l$ and R_m are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{b1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{b2}R_{b3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{b2}R_{b3}$, where R_{b1}, R_{b2} , and R_{b3} are independently selected


from the group consisting of a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof or methacryloyl chloride; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl, where $0 < \alpha < 10$ and $1 < \beta < 25$; a group of aromatic rings having up to 20 carbons in the aromatic framework; fused aromatic rings; CHO; CN; NO₂; Br; Cl; I; phenyl; an acceptor group containing more than two carbon atoms; a functional group derived from an amino acid and $NR_{e1}R_{e2}$; OR_{e3} ; where R_{e1} , R_{e2} , R_{e3} are defined as for R_n and R_o , where R_n and R_o are defined as any member of the group consisting of H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{g1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{g2}R_{g3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{g2}R_{g3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl; aryl groups; fused aromatic rings; polymerizable functionalities;

R_{g1} , R_{g2} , and R_{g3} are independently selected from: H; a linear or branched alkyl group with up to 25 carbons; a functional group derived from an amino acid: a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof or methacryloyl chloride.

5. (Amended) A method for preparing a compound in an electronically excited state, comprising the steps of:

a) exposing a compound having the formula D_1 - Π - D_2 to radiation, wherein D_1 and D_2 are electron donor groups; and Π comprises a bridge of π -conjugated bonds connecting D_1 and D_2 ; and

b) converting said compound to a multi-photon electronically excited state upon simultaneous absorption of at least two photons of said radiation by said compound, wherein the sum of the energies of all of said absorbed photons is greater than or equal to the transition energy from a ground state of said compound to said multi-photon excited state and wherein the energy of each absorbed photon is less than the transition energy between said ground state and the lowest single-photon excited state of said compound and is less than the transition energy between said multi-photon excited state and said ground state [A method according to claim 2], wherein said compound is further defined by a formula



where D_a is selected from the group consisting of N, O, S and P;

where D_b is selected from the group consisting of N, O, S and P;

m, n, o are integers such that $0 \leq m \leq 10$, $0 \leq n \leq 10$, $0 \leq o \leq 10$; and

where:

X, Y, Z are independently selected from the group consisting of: $CR_k=CR_l$; O; S; and N- R_m ;

R_a, R_b, R_c, R_d are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{a1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl; where $0 < \alpha < 10$ and $1 < \beta < 25$, a group of aromatic rings having up to 20 carbons in the aromatic ring framework; fused aromatic rings, vinyl; allyl; 4-styryl; acrolyl; methacroyl; acrylonitrile, isocyanate; isothiocyanate; epoxides; strained ring olefins; $(-CH_2)_\delta SiCl_3$; $(-CH_2)_\delta Si(OCH_2CH_3)_3$; and $(-CH_2)_\delta Si(OCH_3)_3$; where $\delta < 25$;

R_{a1}, R_{a2} , and R_{a3} are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons, a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof, or methacryloyl chloride;

$R_e, R_f, R_g, R_h, R_i, R_j, R_k, R_l$ and R_m are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{b1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{b2}R_{b3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{b2}R_{b3}$, where R_{b1}, R_{b2} , and R_{b3} are independently selected from a functional group derived from an amino acid, a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof or methacryloyl chloride; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl, where $0 < \alpha < 10$ and $1 < \beta < 25$; a group of aromatic rings having up to 20 carbons in the aromatic


framework; fused aromatic rings; CHO; CN; NO₂; Br; Cl; I; phenyl; an acceptor group containing more than two carbon atoms; a functional group derived from an amino acid and NRe₁Re₂; ORe₃; where Re₁, Re₂, Re₃ are defined as for R_n and R_o, where R_n and R_o are defined as any member of the group consisting of H; a linear or branched alkyl group with up to 25 carbons; -(CH₂CH₂O)_α-(CH₂)_βOR_{g1}; -(CH₂CH₂O)_α-(CH₂)_βNR_{g2}R_{g3}; -(CH₂CH₂O)_α-(CH₂)_βCONR_{g2}R_{g3}; -(CH₂CH₂O)_α-(CH₂)_βCN; -(CH₂CH₂O)_α-(CH₂)_βCl; -(CH₂CH₂O)_α-(CH₂)_βBr; -(CH₂CH₂O)_α-(CH₂)_βI; -(CH₂CH₂O)_α-(CH₂)_β-Phenyl; aryl groups; fused aromatic rings; polymerizable functionalities;

R_{g1}, R_{g2}, and R_{g3} are independently selected from: H; a linear or branched alkyl group with up to 25 carbons; a functional group derived from an amino acid: a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof or methacryloyl chloride.

6. (Amended) A method for preparing a compound in an electronically excited state, comprising the steps of:

a) exposing a compound having the formula D₁-II-D₂ to radiation, wherein D₁ and D₂ are electron donor groups; and II comprises a bridge of π-conjugated bonds connecting D₁ and D₂; and

b) converting said compound to a multi-photon electronically excited state upon simultaneous absorption of at least two photons of said radiation by said compound, wherein the sum of the energies of all of said absorbed photons is greater than or equal to the transition energy from a ground state of said compound to said multi-photon excited state and wherein the energy of each absorbed photon is less than the transition energy between said ground state and the lowest single-photon excited state of said compound and is less than the transition energy between said multi-photon excited state and said ground state [A method according to claim 2], wherein said compound is further defined by a formula



where D_a is selected from the group consisting of N, O, S and P;

where D_b is selected from the group consisting of N, O, S and P;

m, n, o are integers such that $0 \leq m \leq 10$, $0 \leq n \leq 10$, $0 \leq o \leq 10$; and where:

X, Y, Z are independently selected from the group consisting of: $CR_k=CR_l$; O; S; and N- R_m ;

R_a , R_b , R_c , R_d are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{a1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl; where $0 < \forall < 10$ and $1 < \exists < 25$, a group of aromatic rings having up to 20 carbons in the aromatic ring framework; fused aromatic rings, vinyl; allyl; 4-styryl; acroyl; methacroyl; acrylonitrile, isocyanate; isothiocyanate; epoxides; strained ring olefins; $(-CH_2)_\delta SiCl_3$; $(-CH_2)_\delta Si(OCH_2CH_3)_3$; and $(-CH_2)_\delta Si(OCH_3)_3$; where $\delta < 25$;

R_{a1} , R_{a2} , and R_{a3} are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons, a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof, or methacryloyl chloride;

R_e , R_f , R_i , R_j , R_k , R_l and R_m are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{b1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{b2}R_{b3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{b2}R_{b3}$, where R_{b1} , R_{b2} , and R_{b3} are independently selected from a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof or methacryloyl chloride; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl, where $0 < \forall < 10$ and $1 < \exists < 25$; a group of aromatic rings having up to 20 carbons in the aromatic framework; fused aromatic rings; CHO; CN; NO₂; Br; Cl; I; phenyl; an acceptor group containing more than two carbon atoms; a functional group derived from an amino acid and $NR_{e1}R_{e2}$; OR_{e3} ; where R_{e1} , R_{e2} , R_{e3} are defined as for R_n and R_o , where R_n and R_o are defined as any member of the group consisting of H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{g1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{g2}R_{g3}$;


$-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{CONR}_{g2}\text{R}_{g3}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{CN}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{Cl}$;
 $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{Br}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{I}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{-Phenyl}$; aryl
 groups; fused aromatic rings; polymerizable functionalities;

R_{g1} , R_{g2} , and R_{g3} are independently selected from: H; a linear or branched alkyl group with up to 25 carbons; a functional group derived from an amino acid: a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof or methacryloyl chloride.

7. (Amended) A method for preparing a compound in an electronically excited state, comprising the steps of:

a) exposing a compound having the formula $\text{D}_1\text{-}\Pi\text{-D}_2$ to radiation, wherein D_1 and D_2 are electron donor groups; and Π comprises a bridge of π -conjugated bonds connecting D_1 and D_2 ; and

b) converting said compound to a multi-photon electronically excited state upon simultaneous absorption of at least two photons of said radiation by said compound, wherein the sum of the energies of all of said absorbed photons is greater than or equal to the transition energy from a ground state of said compound to said multi-photon excited state and wherein the energy of each absorbed photon is less than the transition energy between said ground state and the lowest single-photon excited state of said compound and is less than the transition energy between said multi-photon excited state and said ground state [A method according to claim 2], wherein said compound is further defined by a formula



where D_a is selected from the group consisting of N, O, S and P;

where D_b is selected from the group consisting of N, O, S and P;

m , n , o are integers such that $0 \leq m \leq 10$, $0 \leq n \leq 10$, $0 \leq o \leq 10$; and

where:

X, Y, Z are independently selected from the group consisting of: $\text{CR}_k=\text{CR}_l$; O; S; and N- R_m ;

R_a , R_b , R_c , R_d are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{OR}_{a1}$;

$-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{a2}R_{a3}$;
 $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$;
 $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl; where $0 < \forall < 10$ and $1 < \exists < 25$, a
 group of aromatic rings having up to 20 carbons in the aromatic ring framework; fused
 aromatic rings, vinyl; allyl; 4-styryl; acroyl; methacroyl; acrylonitrile, isocyanate;
 isothiocyanate; epoxides; strained ring olefins; $(-CH_2)_\delta SiCl_3$; $(-CH_2)_\delta Si(OCH_2CH_3)_3$; and
 $(-CH_2)_\delta Si(OCH_3)_3$; where $\delta < 25$;

R_{a1} , R_{a2} , and R_{a3} are independently selected from the group consisting of: H; a
 linear or branched alkyl group with up to 25 carbons, a functional group derived from an
 amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene,
 ruthenocene, cyanuric chloride and derivatives thereof, or methacryloyl chloride;

R_e , R_f , R_i , R_j , R_k , R_l and R_m are independently selected from the group consisting
 of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{b1}$;
 $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{b2}R_{b3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{b2}R_{b3}$, where R_{b1} , R_{b2} , and
 R_{b3} are independently selected from the group consisting of a functional group derived from
 an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene,
 ruthenocene, cyanuric chloride and derivatives thereof or methacryloyl chloride;
 $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$;
 $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl, where $0 < \alpha < 10$ and $1 < \beta < 25$; a
 group of aromatic rings having up to 20 carbons in the aromatic framework; fused aromatic
 rings; CHO; CN; NO₂; Br; Cl; I; phenyl; an acceptor group containing more than two carbon
 atoms; a functional group derived from an amino acid and $NR_{e1}R_{e2}$; OR_{e3} ; where R_{e1} , R_{e2} ,
 R_{e3} are defined as for R_n and R_o , where R_n and R_o are defined as any member of the group
 consisting of H; a linear or branched alkyl group with up to 25 carbons;


$-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{g1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{g2}R_{g3}$;
 $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{g2}R_{g3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$;
 $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl; aryl
 groups; fused aromatic ring; polymerizable functionalities;

R_{g1} , R_{g2} , and R_{g3} are independently selected from: H; a linear or branched alkyl group with
 up to 25 carbons; a functional group derived from an amino acid: a polypeptide; adenine;
 guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and
 derivatives thereof or methacryloyl chloride.


13. (Amended) A method for preparing a compound in an electronically excited state, comprising the steps of:

a) exposing a compound having the formula A_1 - Π - A_2 to radiation, wherein A_1 and A_2 are electron acceptors; and Π comprises a bridge of π -conjugated bonds connecting A_1 and A_2 ; and


b) converting said compound to a multi-photon electronically excited state upon simultaneous absorption of at least two photons of said radiation by said compound, wherein the sum of the energies of all of said absorbed photons is greater than or equal to the transition energy from a ground state of said compound to said multi-photon excited state and wherein the energy of each absorbed photon is less than the transition energy between said ground state and the lowest single-photon excited state of said compound and is less than the transition energy between said multi-photon excited state and said ground state [A method according to claim 9], wherein said compound is further defined by a formula




where A_a and A_b can be independently selected from: CHO; CN; NO₂, and




A1




A2




A3




A4




A5




A6




A7




A8




A9




A10




A11




A12




A13




A14




A15




A16




A17



A18




A19




A20


T00E20"4288T660




A21




A22




A23




A24




A25




A26




A27




A28




A29




A30




A31




A32




A33




A34




A35




A36




A37




A38




A39



A40



A41



A42

in addition A_a and A_b can be independently selected from Br, Cl, and I; and where $0 \leq m \leq 10$, $0 \leq n \leq 10$, $0 \leq o \leq 10$.

; and where:

X, Y, Z are independently selected from the group consisting of: $CR_k=CR_l$; O; S; and N- R_m ;

R_a , R_b , R_c , R_d are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{a1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl; where $0 < \alpha < 10$ and $1 < \beta < 25$, a group of aromatic rings having up to 20 carbons in the aromatic ring framework; fused aromatic rings, vinyl; allyl; 4-styryl; acrolyl; methacroyl; acrylonitrile, isocyanate; isothiocyanate; epoxides; strained ring olefins; $(-CH_2)_\delta SiCl_3$; $(-CH_2)_\delta Si(OCH_2CH_3)_3$; and $(-CH_2)_\delta Si(OCH_3)_3$; where $\delta < 25$;

R_{a1} , R_{a2} , and R_{a3} are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons, a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof, or methacryloyl chloride;

R_e , R_f , R_i , R_j , R_k , R_l and R_m are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{b1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{b2}R_{b3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{b2}R_{b3}$, where R_{b1} , R_{b2} , and R_{b3} are independently selected from the group consisting of a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof or methacryloyl chloride; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl, where $0 < \alpha < 10$ and $1 < \beta < 25$; a group of aromatic rings having up to 20 carbons in the aromatic framework; fused aromatic rings; CHO; CN; NO₂; Br; Cl; I; phenyl; an acceptor group containing more than two carbon atoms; a functional group derived from an amino acid and $NR_{e1}R_{e2}$; OR_{e3} ; where R_{e1} , R_{e2} , R_{e3} are defined as for R_n and R_o , where R_n and R_o are defined as any member of the group consisting of H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{g1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{g2}R_{g3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{g2}R_{g3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$;


$-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{Br}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{I}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta$ -Phenyl; aryl groups; fused aromatic rings; polymerizable functionalities;

R_{g1} , R_{g2} , and R_{g3} are independently selected from: H; a linear or branched alkyl group with up to 25 carbons; a functional group derived from an amino acid; or a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof or methacryloyl chloride.

14. (Amended) A method for preparing a compound in an electronically excited state, comprising the steps of:


a) exposing a compound having the formula A_1 - Π - A_2 to radiation, wherein A_1 and A_2 are electron acceptors; and Π comprises a bridge of π -conjugated bonds connecting A_1 and A_2 ; and

b) converting said compound to a multi-photon electronically excited state upon simultaneous absorption of at least two photons of said radiation by said compound, wherein the sum of the energies of all of said absorbed photons is greater than or equal to the transition energy from a ground state of said compound to said multi-photon excited state and wherein the energy of each absorbed photon is less than the transition energy between said ground state and the lowest single-photon excited state of said compound and is less than the transition energy between said multi-photon excited state and said ground state [A method according to claim 9], wherein said compound is further defined by a formula




where A_a and A_b can be independently selected from: CHO; CN; NO_2 , and


^




A1




A2




A3




A4




A5




A6




A7




A8




A9




A10




A11




A12




A13




A14




A15




A16




A17




A18




A19




A20




A21




A22




A23




A24




A25




A26




A27




A28



A29




A30




A31




A32




A33




A34




A35




A36




A37




A38




A39



A40



A41



A42

in addition A_a and A_b can be independently selected from Br, Cl, and I; and where $0 \leq m \leq 10$, $0 \leq n \leq 10$, $0 \leq o \leq 10$.

; and where:

X, Y, Z are independently selected from the group consisting of: $CR_k=CR_l$; O; S; and N- R_m ;

R_a , R_b , R_c , R_d are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{a1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl; where $0 < \forall < 10$ and $1 < \exists < 25$, a group of aromatic rings having up to 20 carbons in the aromatic ring framework; fused aromatic rings, vinyl; allyl; 4-styryl; acroyl; methacroyl; acrylonitrile, isocyanate; isothiocyanate; epoxides; strained ring olefins; $-(CH_2)_\delta SiCl_3$; $-(CH_2)_\delta Si(OCH_2CH_3)_3$; and $-(CH_2)_\delta Si(OCH_3)_3$; where $\delta < 25$;

R_{a1} , R_{a2} , and R_{a3} are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons, a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof, or methacryloyl chloride;

R_e , R_f , R_i , R_j , R_k , R_l and R_m are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{b1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{b2}R_{b3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{b2}R_{b3}$, where R_{b1} , R_{b2} , and R_{b3} are independently selected from the group consisting of a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof or methacryloyl chloride; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl, where $0 < \forall < 10$ and $1 < \exists < 25$; a group of aromatic rings having up to 20 carbons in the aromatic framework; fused aromatic rings; CHO; CN; NO₂; Br; Cl; I; phenyl; an acceptor group containing more than two carbon atoms; a functional group derived from an amino acid and $NR_{e1}R_{e2}$; OR_{e3} ; where R_{e1} , R_{e2} , R_{e3} are defined as for R_n and R_o , where R_n and R_o are defined as any member of the group consisting of H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{g1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{g2}R_{g3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{g2}R_{g3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$;